

Name : Mark Scheme

Class	Index Number

METHODIST GIRLS' SCHOOL

Founded in 1887



PRELIMINARY EXAMINATION 2023 Secondary 4

Friday
18 August 2023

ADDITIONAL MATHEMATICS Paper 1

4049/01
2 h 15 min

Candidates answer on the Question Paper.
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST


Write your class, index number and name in the spaces at the top of this page.
Write in dark blue or black pen
You may use a HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions.

Give non-exact numerical answers correct to 3 significant figure, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an approved scientific calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.



*Mathematical Formulae***1. ALGEBRA*****Quadratic Equation***

For the quadratic equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$.

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$


Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

1 A calculator must not be used for this question.



In the diagram, AB is parallel to DC , and AC meets BD at X .

Given that $AB = (1 + 3\sqrt{3})$ cm, $CX = (5 - \sqrt{3})$ cm and $CD = (6 + 2\sqrt{3})$ cm, find the exact length of AX . [5]

$$\frac{AX}{5 - \sqrt{3}} = \frac{1 + 3\sqrt{3}}{6 + 2\sqrt{3}}$$

AX

$$\begin{aligned} &= \frac{1 + 3\sqrt{3}}{6 + 2\sqrt{3}} \times (5 - \sqrt{3}) \\ &= \frac{5 - \sqrt{3} + 15\sqrt{3} - 9}{6 + 2\sqrt{3}} \\ &= \frac{14\sqrt{3} - 4}{6 + 2\sqrt{3}} \\ &= \frac{7\sqrt{3} - 2}{3 + \sqrt{3}} \times \frac{3 - \sqrt{3}}{3 - \sqrt{3}} \\ &= \frac{21\sqrt{3} - 21 - 6 + 2\sqrt{3}}{9 - 3} \\ &= \frac{23\sqrt{3} - 27}{6} \text{ or } \frac{23}{6}\sqrt{3} - \frac{9}{2} \end{aligned}$$

2 A curve has the equation $y = \frac{3-x}{e^{1-2x}}$.

(i) Show that the curve is an increasing function of x for $x < \frac{5}{2}$. [5]

$$\begin{aligned}\frac{dy}{dx} &= \frac{e^{1-2x}(-1) - (3-x)e^{1-2x}(-2)}{(e^{1-2x})^2} \\ &= \frac{-1+6-2x}{e^{1-2x}} \\ &= \frac{5-2x}{e^{1-2x}}\end{aligned}$$

$$\text{For } x < \frac{5}{2}, e^{1-2x} > 0$$

$$x < \frac{5}{2}$$

$$2x - 5 < 0$$

$$5 - 2x > 0$$

$$\therefore \frac{dy}{dx} > 0 \text{ for } x < \frac{5}{2}$$

Hence, curve is an increasing function for $x < \frac{5}{2}$

(ii) Given that y is decreasing at a constant rate of 2 units per second, find the rate of change of x when $x = 0$. [2]

$$\begin{aligned}\frac{dy}{dt} &= \frac{dy}{dx} \times \frac{dx}{dt} \\ -2 &= \frac{5-2x}{e^{1-2x}} \times \frac{dx}{dt} \\ -2 &= \frac{5}{e} \times \frac{dx}{dt} \\ \frac{dx}{dt} &= \frac{-2e}{5} = -1.09(3\text{sf})\end{aligned}$$

- 3 The function $f(x) = 2x^3 + ax^2 + bx + 6$, where a and b are constants, is exactly divisible by $x + 2$. When $f(x)$ is divided by $x - 2$, the remainder is -12 .

(i) Find the value of a and of b . [4]

$$\begin{aligned}
 f(-2) &= 0 \\
 -2(-2)^3 + a(-2)^2 + b(-2) + 6 &= 0 \\
 4a - 2b &= 10 \\
 b &= 2a - 5 \quad \text{----- (1)} \\
 f(2) &= -12 \\
 2(2)^3 + a(2)^2 + b(2) + 6 &= -12 \\
 4a + 2b &= -34 \\
 2a + b &= -17 \quad \text{----- (2)} \\
 (2) \text{ into } (1), \\
 2a + 2a - 5 &= -17 \\
 4a &= -12 \\
 a &= -3 \\
 b &= -11
 \end{aligned}$$

(ii) Factorise $f(x)$ completely and hence solve $f(x) = 0$. [3]

$$\begin{aligned}
 2x^3 - 3x^2 - 11x + 6 &= (x + 2)(2x^2 + px + 3) \\
 \text{compare coeff of } x^2, \\
 p + 4 &= -3 \\
 p &= -7 \\
 (x + 2)(2x^2 - 7x + 3) &= 0 \\
 (x + 2)(2x - 1)(x - 3) &= 0 \\
 x &= -2, \frac{1}{2} \text{ or } 3
 \end{aligned}$$


4 It is given that $y = 1 - 3\sin 4x$, for $0^\circ \leq x \leq 180^\circ$.

(i) State the amplitude and period of y . [2]

Amplitude = 3

Period = 90°

(ii) Sketch the graph of $y = 1 - 3\sin 4x$. [3]



(iii) Hence, state the number of solutions for $3\sin 4x = 2$ where $0^\circ \leq x \leq 180^\circ$. [1]

- 5 (a) The equation of a curve is $y = \frac{1}{3} \ln(px + 3)$, where p is a constant to be determined. The gradient of the tangent to the curve at $x = -\frac{1}{2}$ is parallel to $3y = x$. Find the equation of the normal to the curve at $x = -\frac{1}{2}$. [4]

$$\frac{dy}{dx} = \frac{1}{3} \left(\frac{1}{px+3} \right) p$$

$$\frac{p}{3 \left(-\frac{1}{2}p + 3 \right)} = \frac{1}{3}$$

$$p = -\frac{1}{2}p + 3$$

$$\frac{3}{2}p = 3$$

$$p = 2$$

$$x = -\frac{1}{2}, y = \frac{1}{3} \ln 2$$

$$\text{grad of normal} = -3$$

$$y - \frac{1}{3} \ln 2 = -3 \left(x + \frac{1}{2} \right)$$

$$y = -3x - \frac{3}{2} + \frac{1}{3} \ln 2$$

- (b) (i) Express $-2x^2 + 4x - 5$ in the form $a(x + p)^2 + q$. [2]

$$-2x^2 + 4x - 5$$

$$= -2 \left(x^2 - 2x + \frac{5}{2} \right)$$

$$= -2 \left[(x-1)^2 - 1 + \frac{5}{2} \right]$$

$$= -2(x-1)^2 - 3$$

- (ii) Hence, determine, with explanation, if the graph of $y = -2x^2 + 4x - 5$ will intersect the x -axis. [2]

Since coefficient of $x^2 < 0$, graph has maximum turning point at $(1, -3)$.

Maximum value of y is -3 which is less than 0 .

OR The maximum point is below the x -axis.

Hence, graph does not intersect x -axis.

- 6 (a) In the expansion of $(2+x)^n$, where $n > 0$, the coefficient of x^2 is twice the coefficient of x . Find the value of n . [4]

$$(2+x)^n = 2^n + \binom{n}{1} 2^{n-1} x + \binom{n}{2} 2^{n-2} x^2 + \dots$$

$$\binom{n}{2} 2^{n-2} = 2 \binom{n}{1} 2^{n-1}$$

$$\frac{n(n-1)}{2} (2^{n-2}) = 2n(2^{n-1})$$

$$n^2 - n = \frac{4n(2^{n-1})}{2^{n-2}}$$

$$n^2 - n = 8n$$

$$n^2 - 9n = 0$$

$$n(n-9) = 0$$

$$n = 0 \quad \text{or} \quad n = 9$$

$$(NA)$$

- (b) Find the value of the term that is independent of x in the expansion of $\left(2x - \frac{1}{4x^4}\right)^{15}$. [3]

general term

$$= \binom{15}{r} (2x)^{15-r} \left(-\frac{1}{4x^4}\right)^r$$

$$= \binom{15}{r} 2^{15-r} x^{15-r} \left(-\frac{1}{4}\right)^r x^{-4r}$$

$$= \binom{15}{r} 2^{15-r} \left(-\frac{1}{4}\right)^r x^{15-5r}$$

$$15 - 5r = 0$$

$$r = 3$$

$$\text{value} = \binom{15}{3} 2^{12} \left(-\frac{1}{4}\right)^3 = -29120$$

7 It is given that $y = \log_p(px) + 2\log_p(4x-3) - 1$, where p is a positive integer.

(i) Write down the values of x for which y is defined. [1]

$$x > \frac{3}{4}$$


(ii) Show that y can be written as $\log_p(16x^3 - 24x^2 + 9x)$. [3]

$$\begin{aligned} y &= \log_p(px) + 2\log_p(4x-3) - 1 \\ &= \log_p(px) + \log_p(4x-3)^2 - \log_p p \\ &= \log_p \frac{px(4x-3)^2}{p} \\ &= \log_p [x(16x^2 - 24x + 9)] \\ &= \log_p(16x^3 - 24x^2 + 9x) \end{aligned}$$

(iii) Find the value of x for which $y = \frac{1}{\log_{9x} p}$. [3]

$$\begin{aligned} \log_p(16x^3 - 24x^2 + 9x) &= \frac{1}{\log_{9x} p} \\ \log_p(16x^3 - 24x^2 + 9x) &= 1 \div \frac{\log_p p}{\log_p 9x} \\ \log_p(16x^3 - 24x^2 + 9x) &= \log_p 9x \\ 16x^3 - 24x^2 + 9x &= 9x \\ 16x^3 - 24x^2 &= 0 \\ 8x^2(2x - 3) &= 0 \\ x = 0(\text{NA}) \quad \text{or} \quad x &= \frac{3}{2} \end{aligned}$$

- 8 The figure shows a circular lake, centre O , of radius 2 km. A duck swims across the lake from A to B at 3 km/h and then walks clockwise around the edge of the lake from B to C at 4 km/h. If angle $BAC = \theta$ radians and the total time taken is T hours,



- (i) show that $T = \frac{1}{3}(4\cos\theta + 3\theta)$, [4]

$$\angle ABC = \frac{\pi}{2} \text{ (angle in semicircle)}$$

$$\cos\theta = \frac{AB}{2(2)}$$

$$AB = 4\cos\theta$$

$$\angle BOC = 2\theta \text{ (angle at centre} = 2 \times \text{angle at circumference)}$$

$$\text{arc } BC = 2(2\theta) = 4\theta$$

time

$$= \frac{4\cos\theta}{3} + \frac{4\theta}{4}$$

$$= \frac{4\cos\theta}{3} + \theta$$

$$= \frac{1}{3}(4\cos\theta + 3\theta)$$

8 (ii) find the maximum value of T .

[5]

$$T = \frac{4}{3} \cos \theta + \theta$$

$$\frac{dT}{d\theta} = \frac{4}{3}(-\sin \theta) + 1$$

$$-\frac{4}{3} \sin \theta + 1 = 0$$

$$\sin \theta = \frac{3}{4}$$

$$\theta = \sin^{-1} \frac{3}{4} = 0.84806$$

$$\frac{d^2\theta}{d\theta^2} = -\frac{4}{3} \cos \theta$$

$$\theta = 0.84806,$$

$$\frac{d^2\theta}{d\theta^2} = -\frac{4}{3} \cos(0.84806) = -0.8819 < 0$$

T is a maximum.

$$\text{Max } T = \frac{4}{3} \cos 0.84806 + 0.84806$$

$$= 1.73 \text{ h (3sf)}$$

- 9 (a) It is given that $f(x) = (2x-3)^5 + 4$.
Find $f'(x)$ and hence, determine the nature of the stationary point for $y = f(x)$.
[5]

$$f'(x) = 5(2x-3)^4(2)$$

$$= 10(2x-3)^4$$

$$10(2x-3)^4 = 0$$

$$x = \frac{3}{2}$$

x	$\left(\frac{3}{2}\right)^{-}$	$\frac{3}{2}$	$\left(\frac{3}{2}\right)^{+}$
$f'(x)$	> 0	0	> 0
sketch of tangent	/	—	/

At $\left(\frac{3}{2}, 4\right)$, it is a point of inflexion.

- 9 (b) The gradient function of a curve varies directly with the square of $(x - 2)$.
 Given that the gradient of the curve at $(1, 9)$ is 6, find the coordinates of the point at which the curve meets the y -axis. [5]

$$\frac{dy}{dx} = k(x - 2)^2$$

$$\text{At } x = 1, \frac{dy}{dx} = 6$$

$$k = 6$$

$$\frac{dy}{dx} = 6(x - 2)^2$$

$$y = \int 6(x - 2)^2 dx$$

$$= \frac{6(x - 2)^3}{3} + c$$

$$= 2(x - 2)^3 + c$$

$$9 = 2(1 - 2)^3 + c$$

$$c = 11$$

$$y = 2(x - 2)^3 + 11$$

$$\text{At } x = 0, y = -5$$

coordinates are $(0, -5)$

- 10 (a) If $\cos 56^\circ = p$, express $\sin 28^\circ$ in terms of p . [2]

$$\cos 56^\circ = p$$

$$\cos 2(28^\circ) = p$$

$$1 - 2\sin^2 28^\circ = p$$

$$\sin^2 28^\circ = \frac{1-p}{2}$$

$$\sin 28^\circ = \sqrt{\frac{1-p}{2}} \quad \text{or} \quad -\sqrt{\frac{1-p}{2}} \quad (\text{NA})$$

- (b) (i) Prove the identity $\frac{\cos x + \sin x}{\cos x - \sin x} - \frac{\cos x - \sin x}{\cos x + \sin x} = 2 \tan 2x$. [4]

$$\begin{aligned} & \frac{\cos x + \sin x}{\cos x - \sin x} - \frac{\cos x - \sin x}{\cos x + \sin x} \\ &= \frac{\cos^2 x + 2\cos x \sin x + \sin^2 x - (\cos^2 x - 2\cos x \sin x + \sin^2 x)}{\cos^2 x - \sin^2 x} \\ &= \frac{2(2\sin x \cos x)}{\cos^2 x - \sin^2 x} \\ &= \frac{2\sin 2x}{\cos 2x} \\ &= 2 \tan 2x \end{aligned}$$

- (ii) Determine the values of x in the interval $180^\circ < x < 360^\circ$ for which the identity is not valid. [2]


$$2x = 90^\circ, 270^\circ, 450^\circ, 630^\circ$$

$$x = 45^\circ, 135^\circ, 225^\circ, 315^\circ$$

$$\text{For } 180^\circ < x < 360^\circ,$$

$$x = 225^\circ, 315^\circ$$

11



In the diagram, O is the centre of the circle. The diameter AB is produced to C and the line CD is the tangent to the circle at D . The line AD is produced to E such that a circle can be drawn passing through B , C , E and D . The foot of the perpendicular from A to CD produced is F .

Prove that

- (i) triangle CDE is isosceles, [4]

Let $\angle ADF = x$

$\angle ADF = \angle ABD = x$ (alt seg thm)

$\angle ADF = \angle CDE = x$ (vert opp $\angle$)

$\angle CED = \angle ABD = x$ (ext $\angle$, cyclicquad)

$\therefore \angle CED = \angle CDE$

$\therefore \triangle CDE$ is isosceles.

- (ii) AE bisects angle FAB . [3]

$\angle ADB = 90^\circ$ (rt $\angle$ in semicircle)

$\angle DAB = 180^\circ - 90^\circ - x = 90^\circ - x$ ($\angle$ sum of Δ)


$\angle AFD = 90^\circ$ (given)

$\angle FAD = 180^\circ - 90^\circ - x = 90^\circ - x$ ($\angle$ sum of Δ)

$\therefore \angle FAD = \angle DAB$

$\therefore AE$ bisects $\angle FAB$.

- 12 In the diagram, the points A , B and C have coordinates $(2, 4)$, $(7, 9)$ and $(8, 2)$ respectively. The point M lies on AC such that the line BMD is perpendicular to AC .



- (i) Show that $AB = BC$. [2]

$$AC = \sqrt{(7-2)^2 + (9-4)^2} = \sqrt{50}$$

$$BC = \sqrt{(7-8)^2 + (9-2)^2} = \sqrt{50}$$

$$\therefore AC = BC \text{ (shown)}$$

- (ii) Find the equation of BD . [3]

$$\text{grad of } AC = \frac{4-2}{2-8} = -\frac{1}{3}$$

$$\text{grad of } BD = 3$$

Eqn of BD is

$$y - 9 = 3(x - 7)$$

$$y = 3x - 21 + 9$$

$$y = 3x - 12$$


- (iii) Given that the ratio of $DM : MB = 2 : 1$, find the coordinates of D . [2]

$$\text{coord of } M = \left(\frac{2+8}{2}, \frac{4+2}{2} \right) = (5, 3)$$

$$x\text{-coord of } D = 5 - 2(2) = 1$$

$$y\text{-coord of } D = 3 - 2(6) = -9$$

$$\text{coord } D \text{ is } (1, -9)$$



OR

$$\text{translation } \overrightarrow{BM} = \begin{pmatrix} -2 \\ -6 \end{pmatrix}$$

$$\text{translation } \overrightarrow{MD} = \begin{pmatrix} -4 \\ -12 \end{pmatrix}$$

$$\overrightarrow{OD} = \overrightarrow{OM} + \overrightarrow{MD}$$

$$= \begin{pmatrix} 5 \\ 3 \end{pmatrix} + \begin{pmatrix} -4 \\ -12 \end{pmatrix}$$

$$= \begin{pmatrix} 1 \\ -9 \end{pmatrix}$$

$$\text{coord } D \text{ is } (1, -9)$$

- (iv) Find the area of quadrilateral $ABCD$. [2]

$$\begin{aligned} & \frac{1}{2} \begin{vmatrix} 2 & 1 & 8 & 7 & 2 \\ 4 & -9 & 2 & 9 & 4 \end{vmatrix} \\ &= \frac{1}{2} [-18 + 2 + 72 + 28 - (4 - 72 + 14 + 18)] \\ &= 60 \text{ units}^2 \end{aligned}$$

End of Paper.

BLANK PAGE

BLANK PAGE

BLANK PAGE