



ST ANDREW'S JUNIOR COLLEGE

JC2 Preliminary Examinations

Higher 2

CANDIDATE
NAME

--

CLASS

1	9	S		
---	---	---	--	--

CHEMISTRY

9729/02

Paper 2 Structured Questions

2 September 2020

2 hours

Candidate answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name and class on all the work you hand in.

Write in dark blue or black pen.

You may use a HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions in the **spaces provided** on the Question Paper.

The use of an approved scientific calculator is expected, where appropriate.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
Q1		15
Q2		9
Q3		15
Q4		19
Q5		17
Total		75

This document consists of **25** printed pages.

[Turn over


Answer **all** questions in the spaces provided.

- 1 (a) (i) On the axis below, sketch a graph to show the eleven ionisation energies for sodium.

Explain the following features shown in your graph

- the general trend in ionisation energy
- significant change in values between ionisation energies

ionisation energy



.....

.....

.....

.....

.....

.....

[3]

- (ii) State one similarity and one difference (other than difference in energy) between the lowest and highest energy occupied orbital in sodium.

Similarity	
Difference	

[1]


- (b) (i) State two factors that affect the behaviour of beams of positively charged particles in an electric field.

.....

.....


[1]

- (ii) On the diagram below, show how a beam of particles containing sodium ions, copper atoms and electrons deflects as they pass through an electric field at the same speed. Label your answer clearly.




[2]

- (c) Sodium and tert-butanol reacts to give sodium tert-butoxide and hydrogen gas.



- (i) Complete the energy cycle below to calculate the enthalpy change of this reaction, ΔH_{r} .

Your cycle should include relevant data from the *Data Booklet*.



[3]

- (ii) The standard enthalpy change of vapourisation of butanol, tert-butanol and butanone are given below

compound	$\Delta H_{\text{vap}}^\ominus / \text{kJ mol}^{-1}$
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	+46
$(\text{CH}_3)_3\text{COH}$	+44
$\text{CH}_3\text{CH}_2\text{COCH}_3$	+32

By considering the type of interactions involved, suggest reasons for the

- difference in $\Delta H_{\text{vap}}^\ominus$ between tert-butanol and butanone, and
- similarity in $\Delta H_{\text{vap}}^\ominus$ between butanol and tert-butanol.

.....

.....

.....

.....

.....

.....

.....

.....

.....

[3]

- (d) Compound **A** is a constitutional isomer of tert-butanol, $(\text{CH}_3)_3\text{COH}$. **A** exhibits stereoisomerism. It reacts with chlorine in the presence of UV light to form a mono-chlorinated compound **B**, which has four stereoisomers. Draw the four stereoisomers of **B**.

[2]

[Total: 15]

[Turn over

BLANK PAGE

- 2 (a) Explain why benzene prefers to undergo substitution reactions rather than addition reactions.

.....

.....

..... [1]

- (b) The reaction of phenylamine with nitrosonium ion, NO^+ , produces benzenediazonium ion, which is a very useful intermediate in the synthesis of many aromatic compounds.

The mechanism is shown in Fig. 2.1.




Fig. 2.1

- (i) The electrophile, nitrosonium ion, is formed from the reaction between nitrite, NO_2^- , and acid, H^+ .

Write an ionic equation for this reaction.

..... [1]

- (ii) By considering the bonds formed and broken, show the movement of electron pairs using curly arrows for steps 1 and 3 on **Fig. 2.1**. [2]
- (iii) Using appropriate data from the *Data Booklet*, estimate the bond energy of the π bond in $\text{N}=\text{N}$. [1]


- (iv) Given that the bond energy of the π bond in $\text{N}=\text{O}$ is $+405 \text{ kJ mol}^{-1}$, use your answer in (iii) and relevant data from the *Data Booklet* to calculate the enthalpy change of reaction for step 2.

Hence, by considering the effect on the equilibrium, suggest whether a high or low temperature would favour the proton transfer in step 2.

.....
.....
.....

[2]

(c) Benzenediazonium ion reacts with water to form phenol readily.



- (i) Deduce whether water is acting as an electrophile or nucleophile in the reaction. Explain your answer.

.....

 [1]

- (ii) The reaction of benzenediazonium ion with KI(aq) forms phenol and another carbon-containing side product. Name this carbon-containing side product.

..... [1]

[Total: 9]

- 3 Benzyl bromide, $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$, is a strong irritant that affects skin and mucous membranes. It exists as a colourless liquid under standard conditions and is used in military chemical warfare training due to its irritating but non-lethal nature.

(a) (i) The standard enthalpy change of formation of $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$ is $+22.5 \text{ kJ mol}^{-1}$.

Write an equation that illustrates the standard enthalpy change of formation of $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$.

..... [1]

(ii) Predict the sign of ΔS for the reaction in (a)(i).


.....
.....
..... [1]

(iii) Hence, predict whether the formation of $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$ is spontaneous under standard conditions.

.....
.....
..... [1]

- (b) For safety reasons, benzyl bromide has to be hydrolysed before disposal.

The hydrolysis of benzyl bromide is shown in the equation below.



In the first experiment, 25 cm³ of 0.1 mol dm⁻³ sodium hydroxide solution was added to 25 cm³ of 0.0050 mol dm⁻³ of benzyl bromide solution.

Fig. 3.1 shows the concentration of C₆H₅CH₂OH produced against time for this experiment.




Fig. 3.1

The experiment was repeated to find the order of reaction with respect to sodium hydroxide. The total volume was kept constant and the concentration of benzyl bromide was kept the same. The concentration of sodium hydroxide used was 0.20 mol dm⁻³.

The same graph as Fig. 3.1 was obtained in this second experiment.

- (i) Using Fig. 3.1 and the information given, show that the hydrolysis reaction has the following rate equation:

$$\text{rate} = k[\text{benzyl bromide}]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

[3]

- (ii) Draw the mechanism of the hydrolysis reaction between benzyl bromide and hydroxide ion.

[2]

- (iii) Hence, draw a fully labelled reaction pathway diagram for the hydrolysis of benzyl bromide.

[2]

- (iv) Unlike most straight chain primary bromoalkanes, such as 1-bromoheptane, the hydrolysis of benzyl bromide is zero order with respect to hydroxide ions.

Suggest **two** reasons why benzyl bromide undergoes the mechanism in (b)(ii).

.....

.....

.....

.....

.....

.....

[2]

In the third experiment, the same experiment was conducted using 25 cm^3 of 0.1 mol dm^{-3} sodium hydroxide solution and 25 cm^3 of $0.0050 \text{ mol dm}^{-3}$ of benzyl chloride solution.

- (v) On Fig. 3.1, sketch how the graph would look like when $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ was used instead of $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$.

Explain your answer.

.....

.....

.....

.....

.....

..... [3]

[Total: 15]

- 4 Calcium carbonate, CaCO_3 , is one of nature's most abundant raw materials. It is found as a component of seashells, eggshells, corals and pearls. In terms of applications, it has been used in cave paintings, paper, plastics and construction.

(a) (i) Explain why Ca^{2+} has a larger ionic radius than Na^+ .


.....
.....
..... [1]

(ii) Despite Ca^{2+} having a larger ionic radius than Na^+ , the magnitude of the lattice energy of calcium carbonate is still larger than that of sodium carbonate.

Explain why this is so.

.....
.....
..... [1]

Limestone is a type of calcium carbonate-containing rock found naturally. In industry, large quantities of limestone are heated to form calcium oxide. The equation for this thermal decomposition is as shown.



- (b) (i)** The heating of calcium carbonate takes place over a range of temperatures from 1200 K to 1500 K.

With the aid of a Boltzmann distribution diagram, explain the effect of a high temperature range on the rate of thermal decomposition of calcium carbonate.

.....

.....

.....

.....

.....

.....

[3]

- (ii) Under the influence of the calcium cation, the carbonate ion in calcium carbonate undergoes thermal decomposition to form oxide and carbon dioxide in one step. In this mechanism, a σ bond breaks and a π bond forms simultaneously.

Draw the mechanism for the thermal decomposition of the carbonate ion. Show all charges and relevant lone pairs and show the movement of electron pairs by using curly arrows.

[2]

- (iii) The thermal decomposition of both calcium carbonate and calcium ethanedioate, CaC_2O_4 , involve the weakening and breaking of the same bond. Despite this, the temperature for thermal decomposition required for both calcium salt is different.

Predict which calcium salt requires a higher temperature for thermal decomposition. Explain your answer by making reference to the bond which is weakened and broken.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[3]

- (iv) Lithium carbonate behaves like calcium carbonate during thermal decomposition. Suggest an explanation for this.

.....

.....

.....

.....





.....

.....

[2]

- (c) The Solvay Process is a continuous process that uses limestone to produce sodium carbonate.

The Solvay process comprises of the following key steps.



The Solvay Process operates as two cycles, an ammonia cycle and a carbon dioxide cycle, where both ammonia and carbon dioxide can be recycled.

- (i) With reference to the above equations, explain how ammonia and carbon dioxide are recycled.

.....

 [1]

- (ii) Give one advantage and one disadvantage of recycling these gases.

.....

 [2]

NaHCO_3 is soluble in water. However, the use of high concentration of sodium chloride results in the precipitation of NaHCO_3 in step II.

- (iii) State Le Chatelier's Principle and explain how it can be used to explain why NaHCO_3 precipitates in step II.

.....

.....

.....

.....

.....

.....

.....

[2]

In step III, sodium hydrogen carbonate is heated to obtain sodium carbonate.

- (iv) Suggest an experimental method to confirm the completion of this reaction.

.....

.....

.....

[1]

- (v) The sodium carbonate is eventually passed through a dryer and collected as crystals. Suggest a physical method to determine the purity of the sodium carbonate crystals.

.....

.....

.....

[1]

[Total: 19]

- 5 Copper is used globally in electrical wiring, printed circuit boards, generators and electric motors. Copper is also used in cars in various electronic devices, such as batteries and motors.

(a) State the electronic configuration of Cu^+ .

$1s^2$ [1]

(b) The following electrochemical cell was set up.




Fig. 5.1

(i) Construct a balanced equation for the cell reaction and use the *Data Booklet* to calculate the $E^\ominus_{\text{cell}}$ of this cell.

.....

 [1]

(ii) Draw on Fig. 5.1 the direction of electron flow when the reaction in (b)(i) is occurring.

[1]

(iii) Using your answer in (b)(i), calculate $\Delta G^\ominus$ for the reaction.

[1]

- (iv) Predict how the voltage of this cell would change, if at all, when a small amount of aqueous silver nitrate was accidentally spilled into the Cl_2/Cl^- half-cell.

.....

.....

.....

.....

.....

..... [2]

- (c) A car motor is electroplated with copper to replace its worn copper coating. The total surface area to be electroplated is $0.1\text{m} \times 0.2\text{m}$.

A current of 5.0A is used to electroplate one side of the car motor with new copper for 2h . The electrolyte is copper (II) sulfate.

- (i) Calculate the number of copper atoms electroplated on the car motor.

[2]

- (ii) Assume that each copper atom occupies a cube of length $2.98 \times 10^{-12}\text{m}$ and that there is a uniform coverage.

Calculate the depth of copper electroplated in metres, m.

[1]

Copper (II) iodate, $\text{Cu}(\text{IO}_3)_2$, ($M_r = 413.5$) is sparingly soluble in water.

The solubility of copper (II) iodate in water is 1.36 g dm^{-3} .

(d) (i) Calculate the solubility product, K_{sp} , of copper (II) iodate, stating its units

[2]

(ii) Copper (II) bromide is a greyish solid that is highly soluble in water.

Copper (II) bromide solid was added to 500 cm^3 of $0.100 \text{ mol dm}^{-3}$ potassium iodate solution. Determine minimum mass of copper (II) bromide solid that can be added to the solution for a precipitate to form.

[2]

- (e) (i) Write an equation which describes the standard enthalpy change of hydration of $\text{Cu}^{2+}(\text{g})$.

..... [1]

- (ii) Hence, by considering the type of interactions involved, explain why the standard enthalpy change of hydration is always negative.

Explain how the interactions arise.

.....
.....
.....
.....
.....
..... [2]

- (iii) The following data is useful in this question

enthalpy change of hydration of $\text{Cu}^{2+}(\text{g})$ = $-1650 \text{ kJ mol}^{-1}$

enthalpy change of hydration of iodate ions (g) = -460 kJ mol^{-1}

enthalpy change of solution of copper (II) iodate = $-16.2 \text{ kJ mol}^{-1}$

Calculate the lattice energy of copper (II) iodate using data from the above list.

[1]

[Total: 17]