


NAME		Class	
------	--	-------	--

ST ANDREW'S JUNIOR COLLEGE



JC2 PRELIMINARY EXAMINATIONS

Chemistry (9729)

14 September 2018

Paper 3 Free Response

2 hours

Additional Materials: Data Booklet, Writing Paper

READ THESE INSTRUCTIONS:

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Section A

Answer **all** questions. Marks **[60]**

Section B

Answer **one** question. Marks **[20]**

The use of an approved scientific calculator is expected, where appropriate.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **16** printed pages including this page.

Section A

Answer **all** the questions in this section.


- 1 Electrosynthesis in organic chemistry is the synthesis of compounds in an electrolytic cell using inert electrodes. One important process is the Kolbe reaction. The Kolbe reaction is the decarboxylative dimerization of two carboxylate ions. Alkanes can be formed at the anode by the following reaction.



(a) Explain why alkanes are generally unreactive. [1]

(b) (i) Calculate the volume of carbon dioxide produced when a steady current of 5A is passed through the electrolyte for 32 minutes at standard temperature and pressure. [2]

(ii) Methanoate ions undergo the Kolbe reaction to form only 2 gases. Using the axes given below, sketch the variation of pV/RT against p for one mole of **each** gas at the same temperature. Briefly explain your answer. [2]



- 1 (c) The cathodic reaction for the Kolbe reaction can be represented by the following half equation: $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2 + 2\text{OH}^-$.

Write an overall equation for the Kolbe reaction involving ethanoate ions. Hence, calculate $\Delta G^\ominus$ for the Kolbe reaction given that the $E^\ominus_{\text{cell}}$ value for the reaction is +2.28 V.

[2]

- (d) The reaction mechanism of the Kolbe reaction involves a 4-step reaction mechanism as described below:

Step 1: Dissociation of carboxylic acid to form a carboxylate ion.

Step 2: Oxidation of carboxylate ion to form a carboxyl radical, $\text{RCOO}\bullet$.

Step 3: Decomposition of the carboxyl radical to form carbon dioxide gas and an alkyl radical, $\text{R}\bullet$.


Step 4: Formation of a covalent bond between two alkyl radicals.

- (i) Using ethanoic acid as a starting reagent, suggest suitable equations for **Steps 1, 2** and **4** on how ethane can be produced using the Kolbe reaction.

[3]

- (ii) The decomposition of the carboxyl radical to form carbon dioxide gas and an alkyl radical, $\text{R}\bullet$, in **Step 3** can be represented by the following equation: $\text{RCOO}\bullet \rightarrow \text{CO}_2 + \text{R}\bullet$

Outline the mechanism of this step by copying the diagram below and include relevant curly half arrows.




[1]


- (e) In another separate experiment, a new alkane **A**, C_6H_{12} , was produced. When reacted with bromine under ultraviolet light, **A** produced only **one isomeric monobromo** compound, **B**, which does not have a chiral centre. Draw the skeletal formulae of **A** and **B**.

[2]

- 1 (f) (i) Using monohalogenoethane, C_2H_5X , as examples, describe and explain the relative reactivities of chloro- and bromo-compounds in hydrolysis reactions. [2]
- (ii) Halogenoalkanes such as chlorofluoroalkanes, CFCs, were once used as refrigerant fluids and aerosol propellants. In many applications, they have now been replaced by alkanes. This is because CFCs contribute to the destruction of the ozone layer. Explain how CFCs destroy the ozone layer and suggest one potential hazard of using alkanes instead of CFCs. [2]
- (iii) Halogenoalkanes can react with lithium to give organolithium compounds.



These organolithium compounds can react with carbonyl compounds to form alcohols.



(R is alkyl, R' and R'' are either alkyl or H)


Using the above reaction sequence, deduce the structure of a suitable bromoalkane, R-Br, and a suitable carbonyl compound to synthesise hexan-2-ol. [2]

[Total: 19]

- 2 Wastewater has to be treated carefully to remove all harmful contaminants before it can be released into water bodies.

(a) Industrial wastewater contains high levels of heavy metal ions. Electrocoagulation is used to remove heavy metal ions from wastewater. In this method, heavy metal ions are oxidised to form hydroxides, so that they can coagulate easily and be removed easily in subsequent steps.

A sample of wastewater containing Fe^{2+} is treated by electrocoagulation.




- (i) By considering the interaction it forms with water, explain why Fe^{2+} cannot be easily removed from wastewater. [1]
 - (ii) State a suitable physical method to remove $\text{Fe}(\text{OH})_3$ from the wastewater. [1]
 - (iii) Using *Le Chatelier's Principle*, suggest why the wastewater has to be kept alkaline for a more effective removal of Fe^{2+} . [1]
 - (iv) With an appropriate sketch of the Boltzmann distribution, explain how an increase in temperature would affect the rate of the electrocoagulation process. [3]
- (b) Ammonia is a toxic substance present in wastewater from the mining industry. Nitrification is a common biological treatment method to convert ammonia into less toxic nitrate.




- (i) Write the expression for the equilibrium constant, K_c , for this reaction given that K_c has units of $\text{mol}^{-1} \text{dm}^3$. [1]
- (ii) 12 mol of oxygen gas was pumped into 1000 dm^3 of wastewater containing 5 mol of NH_3 . After nitrification, the resulting treated wastewater had a pH of 4.2.
Calculate the value of K_c . Leave your answer to 3 significant figures. [3]
- (iii) State how the K_c in (ii) would change when the amount of NH_3 in wastewater is increased to 10 mol. [1]
- (iv) The nitrification process was carried out at room temperature. It was observed that the outer surface of the reaction vessel was cold after some time.
By considering the effect of temperature on the equilibrium, suggest and explain how you can change the reaction temperature to increase the conversion of ammonia. [2]

- 2 (b) (v) Construct a suitable energy cycle using the following equations to show how the enthalpy change of reaction, ΔH_r , can be determined.



- (c) Wastewater treatment plants that treat pharmaceutical wastewater produces sludge that contains a high amount of organic substances. These organic substances can be combusted to harness energy for other processes. Tonalide, $\text{C}_{18}\text{H}_{26}\text{O}$, is one such organic compound found commonly in sludge. It has a melting point of 54.5°C .




tonalide

- (i) Write an equation which describes the standard enthalpy change of combustion of tonalide. **[1]**
- (ii) The enthalpy change of combustion of tonalide is -865 kJ mol^{-1} .
Given that the combustion is 70% efficient, determine the amount of energy that can be harnessed from the combustion of sludge, which contains 900 g of tonalide. **[2]**
- (iii) The sign of the entropy change of combustion of tonalide is positive. Briefly explain the effect of temperature on the spontaneity of combustion of tonalide. **[1]**


[Total: 20]

- 3 Geraniol, $C_{10}H_{18}O$, is commonly used in perfumes and food flavourings.




Geraniol



It is produced from geranyl diphosphate as shown in the following equation.



Geranyl diphosphate




- (a) The mechanism of this reaction consists of the following:



- (i) State the mechanism in reaction I. [1]
- (ii) Explain why the mechanism in reaction I takes place in two steps instead of one step. [1]
- (iii) State the role of water in reaction II. [1]

- 3 (a) (iv) The following compound can be synthesised from geraniol.



Propose a reaction scheme for this synthesis.

[4]

- (v) Explain what is meant by constitutional isomers and draw a constitutional isomer of geraniol that will form a yellow precipitate with alkaline aqueous iodine.

[2]

- (b) Geranyl diphosphate can be synthesised using P_4O_{10} .

- (i) Write an equation for the reaction of P_4O_{10} in water and describe the effect of the resulting solution on universal indicator solution.
- (ii) Naturally occurring phosphorus contains the isotopes ^{31}P , ^{32}P and ^{33}P . Explain the term isotopes and calculate the relative atomic mass of phosphorus given that the relative abundance of each isotope is 91.1 %, 7.9% and 1.0 % respectively.


[2]

[2]

- (c) Geraniol is a by-product of the metabolism of sorbic acid, C_6H_8COOH . Sorbic acid and its salt are used to prevent growth of mould and fungi and to regulate the pH in food and drinks. With the aid of relevant equations, explain how the pH of food and drinks can be regulated through the use of sorbic acid and its salt.

[3]

- 3 (d) Another constitutional isomer of geraniol is compound **X**. **X** does not react with 2,4-dinitrophenylhydrazine. **X** produces effervescence when sodium metal is added to it. When hot acidified $\text{K}_2\text{Cr}_2\text{O}_7$ is added to **X**, orange solution remains. On addition of aqueous bromine to **X**, the orange solution decolourises and compound **Y** is formed. **X** reacts with hot acidified KMnO_4 to form compound **Z** only. When **Z** is reacted with excess concentrated H_2SO_4 at 170°C , the following product is obtained.



X, **Y** and **Z** are all able to rotate the plane of polarised light. Suggest the identities of **X**, **Y** and **Z**, explaining your reasoning.

[5]

[Total: 21]


Section B

Answer **one** question in this section.

- 4 Transition elements and its compounds have various applications, such as in materials, catalysis and even in nature. An example of a transition element which has many uses and is widely researched on is copper.

(a) Explain what is meant by the term *transition element*. [1]

- (b) (i) Plastocyanin is a copper-containing protein found in vascular plants. The shape around Cu^{2+} in plastocyanin is tetrahedral and can be represented by the following simplified structure. The amino acids found around Cu^{2+} act as ligands.



Define the term *ligand*. [1]

- (ii) Plastocyanin is commonly involved in electron transfer and this results in the formation of Cu^+ complex which has a deep blue colour.

State the electronic configuration of Cu^+ . [1]

- (iii) The deep blue colour seen is considered to be unusual.

Suggest a reason why this is so. [1]

- (c) Azurite is a copper containing mineral that is deep blue in colour. The formula of azurite is $\text{Cu}_3(\text{CO}_3)_2(\text{OH})_2$ and it can be inferred that azurite contained copper(II) carbonate and copper(II) hydroxide, which are sparingly soluble salts.

- (i) To prepare small crystals of azurite in the lab, a few drops of copper(II) sulfate solution can be added into a saturated solution of sodium carbonate and leave to stand overnight.

With reference to the preparation method as described, write an equation to suggest how OH^- may be produced from a saturated solution of sodium carbonate. [1]

- (ii) A student wishes to determine the percentage of azurite in a rock sample. He proposes to do so by performing a direct titration with dilute sulfuric acid.


Suggest an explanation why the proposed method is not ideal. [1]

- 4 (c) (iii) Another student proposed the following method which is ideal in determining the percentage of azurite in a rock sample.

Step 1: Add 7.00 g of the rock sample to 130 cm³ of 0.550 mol dm⁻³ sulfuric acid. The resulting solution was topped up to 250 cm³ using deionised water.

Step 2: 25.0 cm³ of the solution required 15.55 cm³ of 0.200 mol dm⁻³ sodium hydroxide for complete reaction.

Azurite reacts with sulfuric acid according to the following equation.



Complete the balancing of the above equation by deducing values for **a**, **b** and **c**. Hence, calculate the percentage of azurite in the rock sample.

[3]

- (iv) Using your knowledge on the thermal decomposition of carbonates and given that Cu(OH)₂ thermally decomposes to form CuO and H₂O, write an equation for the thermal decomposition of azurite.

Using the equation written and appropriate molar mass values, calculate the percentage loss in mass upon strongly heating a sample of azurite until no further changes.

[3]

- (v) In another instance, a student tried to synthesise azurite by adding copper(II) nitrate solution slowly to a solution containing 0.100 mol dm⁻³ sodium carbonate solution and 0.100 mol dm⁻³ potassium hydroxide solution. Calculate the concentration of Cu²⁺ required to begin precipitating CuCO₃ and Cu(OH)₂ respectively. Hence, state which compound precipitates first assuming that there is a negligible change in volume.

$$K_{sp} \text{ of CuCO}_3 = 7.08 \times 10^{-9} \text{ mol}^2 \text{ dm}^{-6}$$

$$K_{sp} \text{ of Cu(OH)}_2 = 4.8 \times 10^{-20} \text{ mol}^3 \text{ dm}^{-9}$$

[3]

- (vi) When sufficient ammonia is added to copper(II) hydroxide precipitate, the precipitate dissolves. Explain this observation with the aid of relevant equations.

[2]

- (d) Copper metal is obtained from copper ore and is widely used in electrical wiring. To prevent short circuit, the metal used must have high purity. Outline how the impure copper ore is purified industrially to remove impurities such as zinc and silver metals.

[3]

[Total: 20]

- 5 Iron is an important transition element and the second most abundant metal in Earth's crust. Iron containing compounds have various uses and the element itself can also be found in a number of proteins within living things. Iron and its compounds have been widely researched on as they are relatively cheap and non-toxic.

(a) Haemoglobin is a protein found in red blood cells. Within haemoglobin, it contains an Fe^{2+} ion which has an octahedral shape as it is surrounded by five nitrogen-containing ligands and one oxygen-containing ligand, which is H_2O in deoxyhaemoglobin and O_2 in oxyhaemoglobin.

- (i) A species with unpaired electrons is paramagnetic and can be attracted by an externally applied magnetic field. On the other hand, a species with no unpaired electrons is diamagnetic and is unaffected by a magnetic field.


Oxyhaemoglobin is found to be diamagnetic. Based on the information provided and using the diagram below, show the electronic distribution of the 3d electrons of Fe^{2+} in oxyhaemoglobin.



[1]


- (ii) In both oxyhaemoglobin and deoxyhaemoglobin, the 3d orbitals are no longer degenerate as they are split into two different energy levels.

Explain this observation. Using the Cartesian axes like those shown below, draw a fully labelled diagram of one of the 3d orbitals which has a lower energy level.



[3]

- 5 (a) (iii) Explain why oxyhaemoglobin is red. [2]
- (iv) The complementary colours are shown using the following colour wheel.




The wavelength of each colour component in visible light is as provided.

Colour	Purple	Blue	Green	Yellow	Orange	Red
Wavelength / nm	380 - 450	450 - 495	495 - 570	570 - 590	590 - 620	620 - 750

Oxyhaemoglobin is bright red while deoxyhaemoglobin is blue. Given that the wavelength of light is inversely proportional to energy, state and explain if water or oxygen causes a larger energy gap between the 3d orbitals.

[2]


- 5 (a) (v) Fe^{2+} complex in oxyhaemoglobin does not have an overall charge. The structure is as shown.




State the charge around the N atom labelled **(1)** and the bond angle around the N atom labelled **(2)**. [2]

- (vi) Besides Fe^{2+} , Fe^{3+} may also exist in haemoglobin. Explain why Fe can have variable oxidation states in haemoglobin. [1]

- 5 (a) (vii) Each polypeptide chain in haemoglobin is made up of more than 100 amino acid residues. The general structure of an amino acid is as shown below.




In each chain, histidine, leucine and glycine amino acids can be found.

Amino acid	R group
Histidine	
Leucine	$-\text{CH}_2\text{CH}(\text{CH}_3)_2$
Glycine	$-\text{H}$

Draw the structural formula of a tripeptide with the sequence histidine-leucine-glycine at pH 2. [2]

- (viii) Haemoglobin can undergo complete hydrolysis non-enzymatically. State the reagents and condition for this to occur. [1]

- 5 (b) To detect for the presence of blood in a crime scene, an organic compound named luminol may be used. Luminol uses iron in haemoglobin to catalyse the decomposition of hydrogen peroxide, which eventually leads to a blue glow to indicate the presence of blood. The synthetic route for luminol is as shown.



- (i) State the reagents and conditions for **steps 1 to 3**. [3]
- (ii) Draw the structures for intermediate products **X** and **Y**. [2]
- (iii) For **Step 4**, $\text{Na}_2\text{S}_2\text{O}_4$ is commonly used as the reagent. State the role of $\text{Na}_2\text{S}_2\text{O}_4$. [1]

[Total: 20]