



Catholic Junior College

JC2 Preliminary Examinations

Higher 1

CANDIDATE
NAME

CLASS

2T

CHEMISTRY

Paper 2 Structured Questions

8873/02

29 August 2025
2 hours

Candidates answer on the Question Paper.
Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name and class on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.

Section A

Answer **all** the questions.

Section B

Answer **one** question.

The use of an approved scientific calculator is expected where appropriate.
You are reminded of the need for good English and clear presentation in your answers.
A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part of the question.

Section A	Q1	5
	Q2	6
	Q3	15
	Q4	16
	Q5	6
	Q6	12
Section B	Q7	20
	Q8	20
Paper 2 (67%)		80
Paper 1 (33%)		30
Percentage		
Grade		

Section A

Answer **all** the questions in this section in the spaces provided.

- 1(a)** The hydrogen carbonate ion, HCO_3^- , has both acidic and basic properties.
State the conjugate acid and conjugate base of HCO_3^- .

Conjugate acid:

Conjugate base: [1]

- (b)** In the human body, it is important for blood to be maintained at a pH between 7.35 and 7.45. The main buffering agent in blood plasma is the $\text{H}_2\text{CO}_3(\text{aq}) / \text{HCO}_3^-(\text{aq})$ system. During a vigorous physical workout, the muscles will produce an acid which will be transported by the blood to the liver to be broken down.

- (i)** Define the term *buffer solution*.

.....
..... [1]

- (ii)** Briefly explain the effect physical workout will have on the pH of blood if there is no buffer present.

.....
.....
.....
..... [2]

- (iii)** Write an equation to explain how the $\text{H}_2\text{CO}_3(\text{aq}) / \text{HCO}_3^-(\text{aq})$ buffer system helps to maintain the pH of blood during a physical workout.

.....
..... [1]

[Total: 5]

- 2 A student used the apparatus shown in Fig. 2.1 to find the enthalpy change of combustion of butan-1-ol, $\Delta H_c(\text{butan-1-ol})$.




Fig. 2.1

temperature of water before heating = 25.0 °C

temperature of water after heating = 83.7 °C

mass of spirit burner and butan-1-ol before heating = 80.44 g


mass of spirit burner and butan-1-ol after heating = 79.70 g

The specific heat capacity of water is 4.18 J g⁻¹ K⁻¹.



- (a) From the data provided above, calculate the number of moles of butan-1-ol that reacted and hence determine the enthalpy change of combustion of butan-1-ol.

[3]



Complete the table 2.1 below by using the bond energies given in the Data Booklet to calculate another value for the standard enthalpy change of combustion of butan-1-ol.

Table 2.1

Bonds broken:			Bonds formed:	
3 C-C	3 x 350			
9 C-H	9 x 410			
1 C-O	1 x 360			
1 O-H	1 x 460			
6 O=O	6 x 496			

[2]

- (c) Suggest a reason for the discrepancy between the two values calculated in (a) and (b).

.....

 [1]

[Total: 6]

3(a) Propanone reacts with iodine in an aqueous solution of sulfuric acid as shown below.



The rate of this reaction was investigated by withdrawing samples from the reaction mixture, 'quenching' by adding the sample to a large volume of ice-cold water, then measuring the amount of iodine in the sample.

(i) Why is it necessary to quench the sample before measuring the amount of iodine?

.....

 [1]

Separate experiments were carried out to determine the initial rates of reaction and the results obtained are given in the table below.

The rate equation for the reaction is

$$\text{Rate} = k[\text{CH}_3\text{COCH}_3][\text{H}_2\text{SO}_4]^n$$

(ii) The volumes of each reagent is proportional to its concentration. Complete the table below with appropriate values.

Expt	Volume of CH_3COCH_3 / cm^3	Volume of H_2SO_4 / cm^3	Volume of I_2 / cm^3	Volume of H_2O / cm^3	Relative initial reaction rate
1	40	20	20	70	4
2	20	20	20		2
3	40	20	10		
4	40	40	20	50	8

[1]

(iii) Determine the order of reaction, n , with respect to H_2SO_4 . Use data in the table to support your answers.

.....

 [2]

- (b) The rate of hydration of ethanal was studied by dissolving ethanal in 625 cm³ of water.

The concentration of ethanal is measured every 2 minutes. The results of the experiment are shown below.


time /min	[ethanal] /mol dm ⁻³
0	0.80
2	0.56
4	0.42
6	0.31
8	0.23
10	0.16
12	0.12
14	0.09
16	0.08

- (i) Using the information given in the question, calculate the mass of ethanal used in this reaction. (M_r of ethanal = 44.0)

[1]

(ii) On the grid provided below, plot a graph of the results.

[3]



- (iii) Use your graph to show that the reaction is first order with respect to ethanal. Justify your answer by showing appropriate construction lines on your graph.

.....
.....[2]

- (iv) Write the rate equation for the reaction.

.....[1]

- (v) The rate of reaction was measured to be $0.18 \text{ mol dm}^{-3} \text{ min}^{-1}$ at $t = 0 \text{ min}$. Determine the rate constant for the reaction and state its units.


[1]

- (c) Explain how the rate of the reaction will be affected when the experiment is conducted at a lower temperature. Illustrate your answer with the aid of an energy distribution diagram.

.....
.....
.....
.....
.....
.....
.....[3]

[Total: 15]

- 4 This question explores four of the many hundreds of compounds present in a cup of coffee. The first and most well-known compound is caffeine. The structure of caffeine is shown below:




- (a) State the functional group that is circled in the molecule of caffeine.

..... [1]

- (b) (i) In the above structure, circle the alkene functional group. [1]
- (ii) Describe, in terms of orbital overlap, the bonding of the two carbon atoms of the C=C bond in caffeine. A clearly labelled diagram may clarify your answer.

[3]

- (c) The second compound in coffee is damascenone and it has the following structure shown below.



Damascenone

Draw the products formed when damascenone is reacted with the following reagents and conditions:


- (i) H_2 with nickel catalyst in the presence of heat

[2]

- (ii) NaBH_4 in methanol

[1]

- (d) The third compound in coffee is the aldehyde shown below.




- (i) Give the IUPAC name of the aldehyde.

..... [1]


- (ii) Draw the product formed when the aldehyde is reacted with hot acidified potassium manganate (VII), KMnO_4 .

[1]

- (e) The fourth compound, 2,3-pentanedione is present in coffee.



2,3-pentanedione can be reacted in the following sequence:



- (i) State the type of reaction that occurred in step 1.


..... [1]

- (ii) State the reagents and conditions required in step 2.

..... [1]

- (iii) Compound **A** can be used to form polymers. Define the term polymer.

.....
 [1]

compound **A**

- (iv) State the type of polymerisation that compound **A** can undergo.

..... [1]

- (v) Draw a repeat unit of a polymer that compound **A** can form.

[1]

- (vi) Explain why the monomer drawn in part (v) can potentially form thermoset polymers.

.....

..... [1]

[Total: 16]

5(a) Explain the decreasing trend in volatility down Group 17.

.....

.....

.....

.....

.....

.....

.....

..... [2]

(b) Explain the decreasing thermal stability of the hydrides of Group 17.
Include data from the *Data Booklet* in your answer.

.....

.....

..... [2]

(c) Write an ionic equation for the reaction when chlorine gas is bubbled into a solution of potassium iodide. State the colour change observed in the solution of potassium iodide.


.....

.....

..... [2]

[Total: 6]

- 6 Vitamin C, also known as ascorbic acid, has the following structure:




To determine the concentration of ascorbic acid in a typical Vitamin C tablet, an experiment known as *iodometric titration* is used. Firstly, ascorbic acid is reacted with a known amount of iodine that is present in excess. Ascorbic acid will reduce the brown iodine present to colourless iodide. Secondly, unreacted excess iodine is then titrated against a standard solution of sodium thiosulfate. Once the amount of sodium thiosulfate used is known, the amount of ascorbic acid present originally can then be determined.

To prepare a solution of ascorbic acid for the experiment, one 500 mg Vitamin C tablet is dissolved in sulfuric acid and diluted with deionised water up to 250 cm³ mark in a standard flask. 25.0 cm³ of this ascorbic acid solution is pipetted into a conical flask containing 3.00 × 10⁻⁴ mol of iodine solution (excess). Lastly, the resulting solution in the conical flask is immediately titrated with a standard sodium thiosulfate solution using a suitable indicator.


Reaction 1:

Iodine will oxidise ascorbic acid, C₆H₈O₆, to form dehydroascorbic acid:



Reaction 2:

The excess iodine will react with a standardised sodium thiosulfate solution.



It was found 12 cm³ of 0.005 mol dm⁻³ sodium thiosulfate is required to react with iodine.

- (a) With reference to the molecular structure of ascorbic acid, explain why ascorbic acid dissolves in water.

.....
..... [1]

- (b) Explain why **Reaction 2** is a redox reaction. Include oxidation numbers in your answer.

.....
.....
..... [2]

- (c) (i) Calculate the amount of iodine that reacted with sodium thiosulfate.

[2]

- (ii) Calculate the amount of iodine that reacted with 25 cm³ of ascorbic acid solution.

[1]

- (iii) Calculate the amount of ascorbic acid in the 250 cm³ standard solution.

[1]

- (iv) Determine the mass percentage of ascorbic acid in a 500 mg Vitamin C tablet (M_r of ascorbic acid is 176).

[1]

- (d) Suggest a reason why the resulting solution is *immediately* titrated and not allowed to stand for too long before titration.

.....
..... [1]

- (e) Patients who have undergone surgery have been given vitamin C to promote wound healing. Concurrently, external medications with silver ions can also be applied to help prevent inflammation in their post-surgical wounds. Scientists studied the influence of silver nanoparticles on post-surgical wound healing and found that silver nanoparticles exhibited a much more positive effect on wound healing compared to conventional silver preparations.

- (i) A silver cation, Ag^+ , has an ionic **radius** of 1.29×10^{-10} m. Assume that the ions are placed side by side to form the diameter of the smallest spherical silver nanoparticle, calculate the minimum number of Ag^+ ions needed to form the nanoparticle.

[1]

- (ii) With reference to the surface to volume ratio, suggest why silver nanoparticles exhibit a much more positive effect on wound healing compared to conventional silver preparations.

.....

.....

.....


..... [2]

[Total:12]

Section B

Answer **one** question in the spaces provided.

- 7 Dinitrogen tetroxide, N_2O_4 is a brownish yellow liquid which vaporises easily to form a pale yellow gas. It is one of the most important rocket propellants ever developed. N_2O_4 forms an equilibrium mixture with nitrogen dioxide, NO_2 , a brown gas, according to the following equation.

**Equilibrium I**

- (a) (i) Draw a dot-and-cross diagram to show the bonding in a molecule of NO_2 . The molecule contains a dative covalent bond. The central atom is nitrogen which has a single non-bonded valence electron. Distinguish carefully between electrons originating from the central atom and those from the two outermost atoms.

[1]

- (ii) State the shape of the NO_2 molecule.

..... [1]

- (b) Calculate a value for the enthalpy change for the forward reaction in **Equilibrium I**. Relevant enthalpy changes are given in table 7.1 below.

Table 7.1

Thermochemical Equation	Enthalpy change / kJ mol^{-1}
$\frac{1}{2} \text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{NO}_2(\text{g})$	+33.85
$\text{N}_2(\text{g}) + 2 \text{O}_2(\text{g}) \rightarrow \text{N}_2\text{O}_4(\text{g})$	+9.66


[2]

- (c) In an experiment, 0.5 mol of N_2O_4 was allowed to reach equilibrium in a reaction vessel at 50 °C at a pressure of 1 atm. When equilibrium was established, it was found that there were 0.35 mol of N_2O_4 .
- (i) Calculate the number of moles of NO_2 at equilibrium, showing your working clearly.

[2]

- (ii) Given that the reaction was carried out in a vessel that was 2 dm³, determine the equilibrium concentrations of N_2O_4 and NO_2 .

[1]

**Equilibrium I**

(iii) Write the K_c expression for **Equilibrium I**.

[1]

(iv) Using the expression in (iii) and your answers in (ii) determine the value of K_c for this reaction at 50 °C. Show your working clearly.

[1]

(v) The initial experiment in (c) was repeated at a higher pressure of 2 atm and 50 °C.

- I Using Le Chatelier's Principle, predict the effect, if any, on the **equilibrium constant** and **equilibrium composition** of **Equilibrium I**; and
- II State any observations made in terms of colour changes.

.....

.....

.....

.....

.....


.....

.....

.....

..... [4]

- (d) NO_2 reacts with oxygen and water to produce nitric acid, HNO_3 , according to the following equation:



- (i) Nitric acid is a strong acid. Define the term *strong acid*.

.....
 [1]

- (ii) Calculate the pH value of a $0.0250 \text{ mol dm}^{-3}$ solution of dilute nitric acid.

[2]

- (iii) A student suggests that “a $0.0250 \text{ mol dm}^{-3}$ solution of both dilute sulfuric (VI) acid, H_2SO_4 , and nitric acid, HNO_3 , will give the same pH value, because both are strong acids”.

Justify, with relevant calculations of pH, if the student is correct to make that statement.

[2]

- (iv) State and explain which indicator would be suitable for the neutralisation of nitric acid and aqueous ammonia.

.....
 [2]

[Total: 20]

- 8(a) (i)** Phosphorus compounds are commonly used in commercial products. One example is calcium phosphide, Ca_3P_2 , which is used in fireworks.

With reference to structure and bonding, explain the difference in melting point between Ca_3P_2 and elemental phosphorus, P.

.....

.....

.....

.....

.....

.....

.....

..... [3]

- (ii)** When beams of Ca^{2+} and P^{3-} are travelling at the same speed, through an electric field of constant strength, the angle of deflection is proportional to their charge/mass ratio.

Given that Ca^{2+} deflects at an angle of $+x^\circ$ (where the positive sign indicates the direction of deflection), calculate the angle at which P^{3-} deflects.

Your answer must be in terms of x and have the appropriate sign to indicate the direction of deflection.

[1]

- (b) (i) When gaseous PCl_5 is heated in a closed container, the following equilibrium is established.



Sketch, on the same axis below, two graphs that show how the rate of forward and reverse reactions vary over time. Label your graphs clearly and indicate t , the time when equilibrium was established.



[1]

- (ii) Describe the reaction of PCl_5 with water.
Include a balanced equation in your answer.

.....

.....

.....


..... [2]

- (iii) PCl_5 dissolves in some polar solvents without reacting with them. Such solutions conduct electricity. This is due to the presence of the two ions $[\text{PCl}_4]^+$ and $[\text{PCl}_6]^-$. Draw the structures and suggest the shapes of **each** of these ions.

Structure of $[\text{PCl}_4]^+$	Structure of $[\text{PCl}_6]^-$
Shape of ion:	Shape of ion:

[4]

- (c) The first ionisation energies for the elements sodium to potassium are shown in the graph below:



- (i) Write an equation to represent the first ionisation energy of phosphorus.

..... [1]

- (ii) Explain why the first ionisation energy of phosphorus is higher than that of
- silicon

.....

 [2]

- sulfur. (Give suitable electronic configurations to support your answer.)




.....

 [2]

- (d) (i) Calculate the pH of $1.50 \times 10^{-3} \text{ mol dm}^{-3}$ phosphoric acid, $\text{H}_3\text{PO}_4 (\text{aq})$. Assume phosphoric acid behaves as a strong tribasic (triprotic) acid.

[2]

In reality, phosphoric acid is a weak tribasic acid. It dissociates into PO_4^{3-} in the three steps shown below.



- (ii) Write an expression for the first acid dissociation constant of phosphoric acid, K_{a1} .

[1]

- (iii) From the information given above in the three steps, state the weakest acid.

..... [1]

[Total: 20]

BLANK