



VICTORIA JUNIOR COLLEGE
JC 2 PRELIMINARY EXAMINATIONS
Higher 2

CANDIDATE
NAME

CT GROUP

CHEMISTRY

9647/02

Paper 2 Structured

16 September 2014

Candidates answer on the Question Paper.

2 hours

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name and CT group on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a soft pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.

A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
Total	


This document consists of **17** printed pages and **1** blank page.

Answer **all** the questions in the space provided.

1 Planning (P)

Aqueous chlorine, Cl_2 , displaces iodine, I_2 , from aqueous potassium iodide. If a solution of chlorine is mixed with an excess of aqueous potassium iodide, iodine is displaced in a 1:1 molar ratio with chlorine.

The concentration of chlorine in the original solution can therefore be calculated from the concentration of the displaced iodine.



- (a) Write an equation for the reaction between aqueous chlorine, and an excess of aqueous potassium iodide.

.....

[1]

- (b) You are to plan an experiment to determine as accurately as possible the concentration of a saturated aqueous solution of chlorine by titration. The approximate solubility of chlorine is 5 g dm^{-3} at 25°C .

You are provided with the following materials:

- saturated aqueous chlorine;
- solid sodium thiosulfate $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$;
- concentrated aqueous potassium iodide. This will be used in excess.
- the apparatus and chemicals normally found in a school or college laboratory.

Your plan should include details of:

- a calculation of the approximate concentration of saturated aqueous chlorine in mol dm^{-3} at 25°C ;
[A.r: Cl, 35.5]
- a detailed description of the method for preparing a solution of aqueous sodium thiosulfate that can be used in the titration. In a titration, it is usual for the titre volume to be approximately equal to that of the volume of solution pipetted. Calculate the mass of sodium thiosulfate, $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$, which will produce a solution suitable for use in this titration. The relevant calculations and reasoning must be shown in full;
[A.r: H, 1.0; O, 16.0; Na, 23.0; S, 32.1]
- a step-by-step description of how you would carry out sufficient titrations using a suitable indicator to allow an accurate end-point to be obtained;
- an outline calculation to show how the results are to be used to determine the accurate concentration of the aqueous chlorine. Assume that the titre volume used is $w \text{ cm}^3$.

.....

.....

.....

.....

(c) State one hazard that must be considered when planning the experiment and describe a precaution that should be taken to keep risks from this hazard to a minimum.

[Total: 12]

- 2 (a) On Planet Uranus, it is postulated that the number of subshells associated with each principal quantum number and the respective energy levels of the subshells are similar to that on Earth. Each orbital contains a maximum of two electrons. However, the number of orbitals that make up a subshell may or may not be identical to that on Earth.

Figure 1 represents the sketch of the successive ionisation energies of **all** the electrons of an element **T** on Planet Uranus.




Figure 1

- (i) By interpreting figure 1, suggest with reasoning, which period of the Periodic Table element **T** belongs to.

.....

.....

.....

.....

- (ii) Using figure 1, deduce the number of 2p orbitals present in element **T**.

.....

.....


.....

.....

(iii) State the full electronic configuration of element T.

..... [3]

(b) The hydrolysis of ester to form acid and alcohol is an equilibrium reaction. Using ethyl ethanoate as an example, the reaction for the hydrolysis is shown below.



Draw two labelled Maxwell–Boltzmann distribution curves to illustrate clearly how the addition of concentrated sulfuric acid and heating of reaction mixture can increase the rate of hydrolysis. (Written explanation is not required.)

Addition of concentrated sulfuric acid:

Heating of reaction mixture:

[3]

- (c) The rate of hydrolysis (in $\text{mol dm}^{-3} \text{ s}^{-1}$) of ethyl ethanoate at 25°C is given by the following equation.

$$\text{rate} = (4.0 \times 10^{-6} [\text{ester}][\text{H}^+]) + (2.5 \times 10^{-1} [\text{ester}][\text{OH}^-])$$

- (i) Calculate the rate of hydrolysis at $\text{pH} = 1.0$ and 13.0 respectively, given $[\text{ester}] = 0.0100 \text{ mol dm}^{-3}$.

$\text{pH } 1.0$


$\text{pH } 13.0$:

- (ii) Comment on your answers in (c)(i) by relating to the rate equation.

.....

[3]


- (d) Tetracycline is a broad-spectrum polyketide antibiotic indicated for use against many bacterial infections. It is commonly used to treat acne today. Compound **M**, a derivative of tetracycline, has the structure shown below.



compound **M**

Draw the structures of the organic products formed when compound **M** is treated with


- (i) $\text{K}_2\text{Cr}_2\text{O}_7(\text{aq})$, $\text{H}_2\text{SO}_4(\text{aq})$, heat

compound **M**

(ii) aqueous bromine, room temperature

(iii) LiAlH_4 in dry ether, followed by addition of water

(iv) Suggest a simple chemical test to distinguish between compound **M** and compound **N**. You should state clearly the reagents and conditions used, and the observations you would expect to make with each compound.

compound **N**

.....

.....



.....

.....

[6]

[Total: 15]

- 3 Pseudohalogens are a class of compounds that show chemical reactivity similar to halogens. An example of a pseudohalogen and its anion is $(\text{SCN})_2$ and SCN^- respectively.




- (a) (i) Draw the shape of the hybrid orbitals around one of the carbon atoms in thiocyanogen.

- (ii) Draw a diagram to illustrate the shape of thiocyanogen, indicating clearly the covalent bonds and lone pairs of electrons.

- (iii) Thiocyanogen can be synthesised by reacting sodium thiocyanate, NaSCN , with chlorine gas. Write an equation for the formation of thiocyanogen.

..... [4]

- (b) Thiocyanogen can react with hydrocarbons such as methylbenzene via free radical substitution.




- (i) Write equations that describe the following steps.

Initiation

Propagation

Termination (Write down all the possible termination steps.)

- (ii) An isomer of the product shown below was also isolated. Suggest a reason for the formation of this isomer.



.....

[4]

- (c) The standard reduction potential of $(\text{SCN})_2$ is given as



With the aid of the *Data Booklet*, predict the reactions, if any, that might occur when the following reagents are mixed. Write a balanced equation for the reaction that occurs.

- (i) $\text{Na}_2\text{S}_2\text{O}_3$ and $(\text{SCN})_2$

- (ii) acidified $\text{K}_2\text{Cr}_2\text{O}_7$ and $(\text{SCN})_2$

[3]

- (d) SCN^- acts as a ligand in the formation of complex ions. The formation of a complex ion from the hydrated metal ion is shown in the following equilibrium:



where m and n are whole numbers.

The equilibrium constant is called the stability constant, K_{stab} . The table below provides information for the formation of two complexes.


Complex ion	M^{m+}	L^-	n	K_{stab}	Colour
Iron(III) thiocyanate	Fe^{3+}	SCN^-	1	90	Red
Cobalt(II) thiocyanate	Co^{2+}	SCN^-	4	100	Blue

- (i) Write an equation for the formation of cobalt(II) thiocyanate and hence write an expression for its equilibrium constant, K_{stab} and state its units.

- (ii) Calculate the concentration of the complex ion formed when 0.0050 mol of solid cobalt(II) chloride was added to a 2 dm^3 solution of 0.95 mol dm^{-3} KSCN solution.


- (iii) In an experiment to study the stoichiometry of cobalt(II) thiocyanate, a solution of $1.0 \times 10^{-3} \text{ mol dm}^{-3}$ cobalt(II) chloride is mixed with different proportions of a solution of $1.0 \times 10^{-3} \text{ mol dm}^{-3}$ KSCN(aq) such that the total volume of each mixture is kept constant at 10 cm^3 . The absorbance due to the complex ion in each mixture is measured with a colorimeter. The absorbance is directly proportional to the concentration of the complex.

In the grid below, sketch the graph you would expect to obtain.



[6]

- (e) KSCN is used to standardise a solution of silver nitrate. A white precipitate of AgSCN ($K_{sp} = 1.1 \times 10^{-12} \text{ mol}^2 \text{ dm}^{-6}$) is formed during the titration. The indicator used in the titration is yellow aqueous iron(III) solution.



In an experiment, 5 cm³ of 5 mol dm⁻³ nitric acid, 25 cm³ of water and 1 cm³ of iron(III) solution are added to 25.0 cm³ of silver nitrate solution in a conical flask. End-point of titration is reached when 15.00 cm³ of 0.100 mol dm⁻³ KSCN solution is added.

- (i) Calculate the concentration of the original silver nitrate solution.
- (ii) Calculate the concentration of Ag⁺(aq) in the solution at the equivalence point.
- (iii) With reference to the table given in (d), explain why iron(III) solution can be used as an indicator and state the observation at the end-point of the titration.

.....


.....

.....

[4]

[Total: 21]

- 4 (a) Haemoglobin is an important protein present in the red blood cell, which is responsible for transport of oxygen in the body. It contains a haem group and a metal centre as shown below. The four N atoms and the Fe atom are planar.




- (i) State the types of hybridisation at N^1 and N^2 , and the nature of the bonding around Fe atom.

Hybridisation at N^1 : Hybridisation at N^2 :

Type of bond between Fe atom and N^1 :

Type of bond between Fe atom and N^2 :

- (ii) One molecule of haemoglobin (Hb) binds up to four oxygen molecules to form oxyhaemoglobin ($Hb(O_2)_4$), as shown in the following equation.



By applying *Le Chatelier's Principle*, explain how oxygen molecules are transported from the lungs to the body tissues.


.....

- (iii) The percentage saturation of haemoglobin is defined as the percentage of haemoglobin present in the blood that has been converted to oxyhaemoglobin.

With reference to equilibrium (I), state and explain how the percentage saturation of haemoglobin changes when temperature increases.

.....

- (iv) The relationship between the percentage saturation of haemoglobin and concentration of oxygen in blood at different temperatures is shown below.



Suggest the optimum temperature at which haemoglobin operates. Explain your answer.

.....

.....

.....

.....

[8]

- (b) (i) Draw the dot-and-cross diagram of ozone, O_3 .

- (ii) Oxygen–oxygen bond lengths of 3 species are given in the table below.

Species	Oxygen–oxygen bond length / nm
O_3	0.128
O_2^{2-}	0.149
O_2	0.120


With the aid of suitable structures, explain why the two oxygen–oxygen bonds in the ozone molecule have the same bond length which are very different from the peroxide ion and oxygen molecule.

[3]

[Total: 11]

- 5 (a) Eugenoxycetic acid is a colourless and odourless compound. It has shown anti-viral and anti-bacterial properties and is therefore used as antioxidant food preservative in food industry.

Eugenoxycetic acid can be synthesised from eugenol found in clove essential oil as shown in the procedure below via S_N2 mechanism.



Preparation of solution A

1. Dissolve 1.0 g of 2-chloroethanoic acid in 5.0 cm³ of distilled water in a beaker. Add Na₂CO₃(s) slowly until the solution just becomes alkaline.

Preparation of eugenoxycetic acid

2. Dissolve 0.6 g of NaOH(s) in 3.0 cm³ of distilled water in a conical flask. Add 2.0 cm³ eugenol to the flask.
3. Add solution A to the flask in step 2 and keep stirring the mixture for 60 min.
4. Cool the reaction mixture to room temperature and add HCl(aq). Collect the solid formed by suction filtration. Wash the solid with water to obtain a pale yellow solid which is the crude product.

- (i) How can you tell that the solution has just turned alkaline in step 1?

.....

.....

- (ii) Explain the role of NaOH(s) in step 2.

.....

.....

.....

- (iii) Write a balanced equation for the reaction in step 3.

- (iv) Comment if an excess amount of NaOH(s) can be used in step 2.

.....

.....

.....

- (v) Give a simple chemical test to confirm that eugenol is absent in the final product.


.....

.....

.....

[6]

- (b) The structure of a small protein is given below, along with the abbreviated names of its constituent amino acids: **val**, **gln** etc.



- (i) Use the abbreviated names of the amino acids to state the primary structure of the above protein.
-
- (ii) This protein is ionised at pH 7. Identify **one** functional group that could be ionised at pH 7 by circling it on the structure above.
- (iii) The biological function of the protein depends on its three dimensional shape. Describe how the particular amino acids in the above protein are likely to be involved in maintaining its three-dimensional shape.

.....

.....

.....

.....

.....

.....

- (iv) The activity of the above protein is destroyed by heating it under reflux with 6 mol dm^{-3} sulfuric acid for several hours.

Draw the structures of the organic products formed under these conditions from the part of the above protein containing the amino acids: **val-gln**.

[7]

[Total: 13]